

Blastomycosis presenting as an isolated progressive painless verrucous skin lesion

Saira Butt 

Division of Infectious Diseases,
Indiana University School of Medicine,
Indianapolis, IN, USA

Correspondence

Saira Butt, Division of Infectious Diseases,
Indiana University School of Medicine,
545 Barnhill Drive, Emerson Hall 421,
Indianapolis, IN 46202, USA.
Email: sairabuttmd@gmail.com

Abstract

A painless progressive verrucous skin plaque in an endemic area should raise suspicion for blastomycosis resulting in prompt biopsy with fungal stains and culture. Skin is a common extrapulmonary site. Itraconazole is treatment of choice.

A 57-year-old Caucasian man from Indiana was referred to infectious diseases clinic for a skin lesion. A nonpruritic and painless skin lesion had been present on his lower back for a year and had progressively enlarged. Topical steroids worsened the lesion. Patient was asymptomatic. Physical examination showed a 10 × 3 cm raised erythematous verrucous plaque with stuck-on appearance with dry scaly raised brown borders.¹ (Figure 1) The patient worked as a car mechanic, lived in a suburban old house, denied any

travel, was monogamous with his wife, and had no adventurous hobbies. His labs and chest X-ray were normal. The lesion was biopsied and showed thick-walled fungal spores in the dermis with broad-based budding yeast consistent with blastomycosis.² Patient's urine and serum blastomyces antigens were negative. Oral itraconazole 200 mg twice a day was initiated. Itraconazole was tolerated well, and patient had adequate drug level. On 3-month follow-up, the lesion had become flat and less scaly. (Figure 2) On 6-month follow-up,



FIGURE 1 Initial presentation: painless progressive verrucous skin lesion



FIGURE 2 Three-month follow-up: flat and less scaly skin lesion

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.



FIGURE 3 Six-month follow-up: smooth and faded skin lesion

it had become smooth and faded. (Figure 3) Itraconazole was stopped at 6 months, and patient continued to do well clinically without any new symptoms or lesions.

CONFLICT OF INTEREST

None declared.

AUTHOR CONTRIBUTIONS

SB: drafted and reviewed the article.

ORCID

Saira Butt  <https://orcid.org/0000-0003-3273-8379>

REFERENCES

1. Saccente M, Woods GL. Clinical and Lab update on Blastomycosis. *Clin Microbiol Rev.* 2010;23(2):367-381.
2. Schwartz J, Salfelder K. Blastomycosis. A review of 152 cases. *Curr Top Pathol.* 1977;65:165-200.

How to cite this article: Butt S. Blastomycosis presenting as an isolated progressive painless verrucous skin lesion. *Clin Case Rep.* 2020;8:778–779. <https://doi.org/10.1002/ccr3.2556>